arXiv:astro-ph/0312320v1 12 Dec 2003

350μm observations of local IRAS galaxies using SHARC-II

Colin Borys

California Institute of Technology, Pasadena CA 91125, USA

1 SEDs and redshifts of sub-mm bright galaxies

The cosmologically significant population of dusty galaxies detected at sub-mm and mm wavelengths by SCUBA and MAMBO have played a central part in many of the discussions at this workshop. Obtaining their redshifts is critical in understanding their role in galaxy formation and evolution, but this is a challenging task. Nevertheless, Chapman et al. [1] have made significant progress by obtaining spectroscopic redshifts of the subset of sub-mm galaxies detected in deep radio images that have optical counterparts. This exploits the well-known Far-Infrared/Radio correlation, and allows the high resolution radio maps to be used as a surrogate for finding the correct optical counterpart for which spectroscopy can then be attempted.

Because of the large investment of telescope time required to obtain and confirm the spectroscopic redshifts, and because the radio-detected sample consists only of the brighter sub-mm objects (only a few percent of the sub-mm source counts), some groups [2,3,4] have developed photometric techniques to estimate the redshifts of SCUBA/MAMBO detected galaxies. These are also potentially biased because the Spectral Energy Distribution (SED) basically follows a modified black-body curve, which results in a fundamental degeneracy between redshift and dust temperature [5]. However, by assuming that high-redshift galaxy SEDs resemble local ones, a library of templates derived from nearby galaxies can be used to essentially “fix” the temperature and break the degeneracy. One problem with this approach is that the SEDs of local infrared-bright systems are poorly constrained for all but a few of brightest local sources.

One way to address this issue is to obtain photometry at different wavelengths. However, the atmosphere is opaque to FIR emission except for the narrow windows at relatively long wavelengths that SCUBA and MAMBO have already exploited. Efficient observations at the shorter wavelength windows at 350 and 450μm have proved very difficult until now.

2 SHARC-II observations of SLUGS galaxies

The CSO recently commissioned the “CCD-style” SHARC-II camera [6] optimized for use at 350μm. The combination of 384 sensitive detectors and a dish with low surface error allow observations previously impractical with other cameras operating at similar wavelengths.

Using SHARC-II, we have been imaging the sample of 106 IRAS selected galaxies in the highly successful SCUBA Local Universe Galaxy Survey (SLUGS) [7]. These sources are lower luminosity analogues to the high-redshift SCUBA population. With observations at 60, 100 (IRAS) and 850μm (SCUBA), the SEDs were fit and dust parameters estimated. SCUBA was only able to detect 17 of these galaxies at 450μm, but from them it was recognized that the models required a component of cold dust to fit the the SED properly [8]. These results highlight the need for more short-wavelength observations. To date, we have imaged 60 of the 106 SLUGS sources with SHARC-II and have and detected every one. We are also conducting complementary observations of the Chapman et al. sample of SCUBA galaxies with spectroscopic redshifts. Using this spec-z to break the T/$(1 + z)$ degeneracy, the combination of SCUBA and SHARC-II observations will allow a study of the physical dust properties as a function of redshift.

References

1. S.C. Chapman, A.W. Blain, R.J. Ivison, I.R. Smail: Nature **422**, 695 (2003)
2. D. Hughes et al.: MNRAS **335**, 871 (2002)
3. I. Aretxaga et al.:MNRAS **342**, 759 (2003)
4. T. Wiklind: ApJ **588**, 736 (2003)
5. A.W. Blain, V. E. Barnard, S.C. Chapman: MNRAS **338**, 733 (2003)
6. C.D. Dowell et al.: SPIE **4855**, 73 (2003)
7. L. Dunne, S.E. Eales, M. Edmunds, R. Ivison, P. Alexander, D. Clements: MNRAS **327**, 697 (2001)
8. L. Dunne, S.E. Eales: MNRAS **327**, 697 (2001)



Fig. 1. Two IRAS galaxies imaged with SHARC-II. Overlaid on DSS images are SHARC-II Signal-to-Noise contours at 4 and 6 sigma/beam. The black outline denotes the area sampled by the SHARC-II array.